www.MedChemExpress.com

MCE®
MedChemExpress

Recombinant Proteins

- High Purity
- Full Range Sizes
- Superior Biological Activity
- GMP-grade Proteins

Recombinant Proteins

MedChemExpress (MCE) provides a comprehensive catalog of recombinant proteins with various tags, multiple species, excellent lot-to-lot consistency, superior biological activity and certified low levels of endotoxin to meet the needs of all types of customers. MCE's recombinant proteins include: cytokines and growth factors, viral proteins, immune checkpoint proteins, CAR-T related proteins, CD antigens, receptor proteins and enzymes. MCE's recombinant proteins have been cited in research articles covering many different fields and disciplines, such as cell growth and differentiation, cell signaling, biopharmaceutical target discovery, protein structural and functional analyses, etc. By ensuring high-quality products and professional pre-sale and after-sale services, MCE is now being regarded as a partner-of-choice by millions of scientists and technicians.

- **Broad Categories:** Cytokines and Growth Factors, Immune Checkpoint Proteins, CAR-T Related Proteins, CD Antigens, Fc Receptor Proteins, Receptor Proteins, Enzymes & Regulators, Complement System Related Proteins, Ubiquitin Related Proteins, Biotinylated Proteins, Viral Proteins, GMP-grade Proteins
- **Low Endotoxin Levels:** Measured by LAL assay
- **High Purity:** Tested by SDS-PAGE & HPLC
- **Superior Biological Activity:** Validated by relevant in vitro or in vivo assays
- **Excellent Lot-to-Lot Consistency:** Confirmed by Lot-to-Lot data
- **Full Range Sizes:** Different pre-packaged sizes for various needs
- **Competitive Price:** High quality with a reasonable price



High Purity

The purity of human BMP-2 is greater than 95% as analyzed by SDS-PAGE under reducing (R) and non-reducing (N) condition.



Superior Biological Activity

Human GM-CSF stimulates cell proliferation of TF-1 cells with an ED_{50} of less than 0.5 ng/mL.



Excellent Lot-to-Lot Consistency

The ED_{50} of MCE human VEGF165 from three different Lots are similar.



Activity Comparison

The ED_{50} of human VEGF165 from MCE's each Lot is lower than of Competitor P.



Publications Citing Use of MCE Products

Nature. 2025 Feb;638(8052):1112-1121.

Nature. 2025 Feb;638(8050):490-498.

Nature. 2025 Feb;638(8050):519-527.

Nature. 2025 Feb;638(8052):1095-1103.

Nature. 2025 Jan;637(8046):726-735.

...

Cell. 2025 Feb 20;188(4):998-1018.e26.

Cell. 2025 Feb 6;188(3):814-831.e21.

Cell. 2025 Feb 6;188(3):653-670.e24.

Cell. 2025 Jan 23;188(2):412-429.e27.

Cell. 2025 Jan 9;188(1):157-174.e22.

...

Science. 2025 Jan 10;387(6730):eads9960.

Science. 2024 Dec 13;386(6727):eado5561.

Science. 2024 Nov 15;386(6723):eadh9215.

Science. 2024 Sep 13;385(6714):eadj1979.

Science. 2024 Aug 30;385(6712):eado6593.

...

MCE Global Partners

Millipore Sigma · Dana-Farber Cancer Institute · Imperial College London · Harvard University · GSK

Karolinska Institutet · NIH National Institutes of Health · Turning Discovery Into Health · Yale University · Takeda · Roche

AstraZeneca · Northwestern University · Stanford University · Lilly · NUS National University of Singapore

Reconstitution and Storage

1 Centrifuge the tube before opening

During shipment, the protein may adhere to the wall or cap of the vial. Before opening the vial, please centrifuge at 10,000-12,000 rpm for 30 seconds to gather the protein at the bottom of the vial. If a high-speed centrifuge is not available, please centrifuge at 3,000-3,500 rpm for 5 mins.

2 After centrifugation, add the reconstitution buffer to the lyophilized protein powder and mix gently by pipetting. Resuspend in the reconstitution buffer to recommended concentration (no less than 100 μg/mL).

Note: Vigorous vortexing should be avoided as it can cause protein foaming and denaturation, thereby affecting the protein activity.

3 Once reconstituted, recombinant proteins can be stored no more than a week at 2-8°C.

For experiments with a short cycle (no more than 7 days), the recombinant protein solution can be directly added to the culture system and used up within a week. If the experimental concentration is lower than the reconstituted concentration, dilution can be done with a solution containing carrier proteins.

4 For long-term storage, the protein solution should be diluted further with carrier proteins (0.1% BSA, 5% HAS, 10% FBS or 5% trehalose), and then aliquot and stored at -20°C to -80°C.

It is not recommended to freeze the reconstituted product directly at -20°C to -80°C. Some recombinant proteins may stick to the plastic tube wall easily, which results in a lower concentration of protein in the solution and ultimately reduces its activity. Carrier proteins can prevent products from sticking to the tube wall by pre-blocking the protein binding site. Therefore, for long-term storage, cytokines or proteins should be further diluted with the solution containing carrier proteins before making aliquots and freezing.

Note: Avoid repeated freeze/thaw cycles. Each freeze/thaw cycle will cause denaturation or conformational changes in some proteins, thereby reducing the binding ability of antibodies and accelerating protein degradation.

CONTENTS

Cytokines and Growth Factors 2
Immune Checkpoint Proteins 4
CAR-T Related Proteins 6
CD Antigens 7
Fc Receptor Proteins 8
Receptor Proteins 9
Enzymes & Regulators 10
Complement System Related Proteins 11
Ubiquitin Related Proteins 12
Viral Proteins 13
Biotinylated Proteins 14
GMP-grade Proteins 15
Animal-free Recombinant Proteins 15
Recombinant Protein Expression Service 16
Customer Validation 17
Publications Citing Use of MCE Recombinant Proteins 18



Cytokines and Growth Factors

Cytokines are a large class of low molecular weight proteins, peptides, or glycoproteins that are secreted by various types of immune cells such as macrophages and lymphocytes, as well as other cell types such as endothelial cells. They play an important role in regulating cell growth, differentiation, and activation and are involved in many aspects of the innate and adaptive immune response[1][2][3]. Growth factors are soluble signaling molecules that stimulate various cellular processes during development and tissue healing, including cell proliferation, migration, differentiation, and multicellular morphogenesis[4][5]. The terms "growth factors" and "cytokines" are often used interchangeably[5][6].

Human IGF-I (HY-P7018)

The purity of human IGF-I is greater than 95% as analyzed by SDS-PAGE under reducing (R) conditions.



Human IGF-I (HY-P7018)

The ED_{50} of human IGF-I is < 5.0 ng/ml as measured by FDC-P1 cells, corresponding to a specific activity of >2.0 × 10^5 units/mg.



Human EGF (HY-P7109)

The purity of human EGF is greater than 95% as analyzed by SDS-PAGE under reducing (R) conditions.



Human EGF (HY-P7109)

The ED_{50} of human EGF is <0.2 ng/mL as measured by murine BALB/c 3T3 cells, corresponding to a specific activity of 5.0 × 10^6 units/mg.



Cat. No.	Product Name	Species	Source	Tag
HY-P7080	IL-4	Mouse	Mammalian Cells	Tag-free
HY-P7223	IL-6R alpha	Human	Mammalian Cells	Tag-free
HY-P7025	IFN-gamma	Human	*E.coli*	Tag-free
HY-P7287	SDF-1 beta/CXCL12	Human	*E.coli*	Tag-free

Cat. No.	Product Name	Species	Source	Tag
HY-P7085A	TNF-alpha/TNFSF2	Human	Mammalian Cells	Tag-free
HY-P7085	M-CSF	Mouse	*E.coli*	Tag-free
HY-P7118	TGF beta 1	Human	Mammalian Cells	Tag-free
HY-P7086	Noggin	Mouse	Mammalian Cells	Tag-free
HY-P7007	BMP-4	Human	*E.coli*	Tag-free
HY-P70593	Fibronectin	Human	*E.coli*	Tag-free
HY-P70453B	Wnt3a Surrogate	Human	Mammalian Cells	Fc
HY-P7109	EGF	Human	*E.coli*	Tag-free
HY-P70311	Activin A	Human/Mouse/Rat	Mammalian Cells	Tag-free
HY-P7319	AITRL/TNFSF18	Mouse	*E.coli*	Tag-free
HY-P75509	Angiopoietin-2	Canine	Mammalian Cells	N-His
HY-P72650	FGF-21	Cynomolgus	Mammalian Cells	C-His
HY-P703928	BMP1	Human	*E.coli*	C-His
HY-P71827	Adiponectin/ADIPOQ	Bovine	*P.pastoris*	N-His
HY-P7008	BMP-7	Human	*E.coli*	Tag-free
HY-P7257	CCL4	Human	*E.coli*	Tag-free
HY-P70450	CCL5	Human	*E.coli*	Tag-free
HY-P7143	CCL6	Mouse	*E.coli*	Tag-free
HY-P7772	CCL9	Mouse	*E.coli*	Tag-free
HY-P70138	DLK-1	Human	Mammalian Cells	C-His
HY-P7004	FGF basic/bFGF	Human	*E.coli*	Tag-free
HY-P7170	FGF-10	Mouse	*E.coli*	Tag-free
HY-P70533	FGF-8	Human	*E.coli*	Tag-free
HY-P7015A	G-CSF	Human	Mammalian Cells	Tag-free
HY-P7016	GM-CSF	Human	Mammalian Cells	Tag-free
HY-P7017	HB-EGF	Human	*E.coli*	Tag-free
HY-P7018	IGF-I	Human	*E.coli*	Tag-free
HY-P7368	IGFBP-2	Human	Mammalian Cells	C-His
HY-P7027	IL-1 alpha	Human	*E.coli*	Tag-free
HY-P7030A	IL-10	Human	Mammalian Cells	Tag-free
HY-P7049	LIF	Human	*E.coli*	Tag-free
HY-P7051A	Noggin	Human	Mammalian Cells	Tag-free
HY-P70781	SCF	Human	*E.coli*	Tag-free
HY-P70467	SHH	Human	*E.coli*	Tag-free

Immune Checkpoint Proteins

Immune checkpoint (ICP) molecules are ligand-receptor pairs that have an inhibitory or stimulatory effect on the immune response. Most of the ICP proteins that have been described are expressed on cells of the adaptive immune system. ICP proteins act as important immune regulators in maintaining immune homeostasis and immune tolerance, and some cancer cells can bind co-inhibitory receptor molecules to limit the normal anti-tumor immune response, thereby assisting immune escape. ICP therapy for cancer includes strategies that target these regulatory pathways to reinvigorate the anti-tumor function of immune cells[7][8].



Figure 1. Common inhibitory and stimulatory immune checkpoint pathways[9]

Cat. No.	Product Name	Species	Source	Tag
HY-P70691	CTLA-4	Human	Mammalian Cells	C-GST
HY-P70632	PD-L1	Mouse	Mammalian Cells	C-His
HY-P7395	PD-1	Human	Mammalian Cells	C-hFc
HY-P70482	TIM3	Human	Mammalian Cells	C-His
HY-P70722	LAG-3	Human	Mammalian Cells	C-His
HY-P70624	TIGIT	Human	Mammalian Cells	C-His
HY-P7327	CD276/B7-H3	Human	Mammalian Cells	C-His
HY-P7446	4-1BBL/TNFSF9	Mouse	Mammalian Cells	N-His
HY-P7144	CD40L/CD154/TRAP	Human	*E.coli*	Tag-free
HY-P70652	CD276/B7-H3	Human	Mammalian Cells	C-His
HY-P7678	BTLA/CD272	Human	Mammalian Cells	C-His
HY-P7685	BTN3A3	Human	Mammalian Cells	C-His
HY-P70535	Galectin-9	Human	Mammalian Cells	C-His
HY-P7366	HVEM	Human	Sf9 insect Cells	C-hFc
HY-P70494	Nectin-1	Human	Mammalian Cells	C-His
HY-P70807	PVR/CD155	Human	Mammalian Cells	C-His
HY-P73370	PD-L2	Rat	Mammalian Cells	C-hFc
HY-P77879	VISTA	Human	Mammalian Cells	C-hFc
HY-P75610	CD137/4-1BB	Canine	Mammalian Cells	C-His
HY-P7394	OX40/TNFRSF4	Human	Mammalian Cells	C-His
HY-P77457	OX40 Ligand/TNFSF4	Cynomolgus	Mammalian Cells	N-mFc
HY-P73499A	CD40	Human	Mammalian Cells	C-His
HY-P73306	Nectin-3	Human	Mammalian Cells	C-His
HY-P71248	PVRIG	Human	Mammalian Cells	C-mFc
HY-P72468	SIRP alpha	Mouse	Mammalian Cells	C-His
HY-P73121	IDO	Human	*E.coli*	Tag-free
HY-P76396	ICOS	Human	Mammalian Cells	C-His-hFc
HY-P76393	ICOSLG	Cynomolgus	Mammalian Cells	C-His
HY-P72353	CD28	Human/Cynomolgus	Mammalian Cells	C-Fc-Avi
HY-P72033	LIGHT	Human	Mammalian Cells	N-hFc-Myc
HY-P73076	GITR	Human	Mammalian Cells	C-His
HY-P7318	GITRL/AITRL	Human	*E.coli*	Tag-free
HY-P72887	CD200	Human	Mammalian Cells	C-hFc
HY-P76780	CD200R1	Cynomolgus	Mammalian Cells	C-His

CAR-T Related Proteins

Chimeric antigen receptor T (CAR-T) cells, genetically engineered to express synthetic chimeric antigen receptors, can specifically target antigens and kill tumor cells[10][11]. CAR-T cells can specifically recognize their target antigens through single-stranded fragment variant (scFv) binding domains, leading to T cells in a major histocompatibility complex (MHC)-independent manner activation. Therefore, in CAR design, antigen selection is critical for killing target tumor cells[12].



Figure 2. A diagram of a CAR[13]

Cat. No.	Product Name	Species	Source	Tag
HY-P7656	BCMA/TNFRSF17	Mouse	Mammalian Cells	C-Fc
HY-P72019	Siglec-2/CD22	Human	Mammalian Cells	C-His
HY-P70505	CD19	Human	Mammalian Cells	C-Fc
HY-P70731	CD38	Human	Mammalian Cells	C-His
HY-P70148	Mesothelin	Human	Mammalian Cells	C-His
HY-P70189	EGFR vIII`	Human	Mammalian Cells	C-His
HY-P70125	CD276/B7-H3	Cynomolgus	Mammalian Cells	C-His
HY-P70155	EpCAM/TROP1	Human	Mammalian Cells	C-Fc
HY-P70759	HER2/CD340	Human	Mammalian Cells	C-Fc
HY-P70301	Mucin-1	Human	Mammalian Cells	C-Fc
HY-P71031	Siglec-6	Human	Mammalian Cells	C-Fc
HY-P70487	Glypican-3/GPC3	Human	Mammalian Cells	C-His
HY-P70296	Folate receptor alpha	Human	Mammalian Cells	C-His
HY-P70497	CD7	Human	Mammalian Cells	C-His

CD Antigens

Cluster of differentiation (CD) antigens are cell surface molecules that can be used to identify and investigate their presence in leukocytes. Some CD antigens frequently act as cell-cell or cell-matrix adhesion molecules, cytokine receptors, ionophores, or nutrient transporters. CD antigens are routinely used as cellular markers that can be used to identify and isolate the presence and proportion of specific leukocyte populations and lymphocyte subpopulations using fluorescently labeled antibodies[14][15].



Figure 3. CD antigens of T cells[16]

Cat. No.	Product Name	Species	Source	Tag
HY-P70090	CD3 epsilon	Cynomolgus	Mammalian Cells	C-Fc
HY-P72702	CD8 beta	Human	Mammalian Cells	N-His
HY-P70486	CD28	Human/Cynomolgus	Mammalian Cells	N-His
HY-P7321	B7-1/CD80	Human	Mammalian Cells	C-hFc
HY-P70029	Basigin/CD147	Human	Mammalian Cells	C-His
HY-P70507	CD44	Human	Mammalian Cells	C-His
HY-P7679	BTLA/CD272	Human	Mammalian Cells	C-Fc
HY-P7780	Nectin-2/CD112	Human	Mammalian Cells	C-His
HY-P7785	CD127/IL-7RA	Human	Mammalian Cells	C-Fc-His
HY-P70550	CD137/4-1BB	Cynomolgus	Mammalian Cells	C-His
HY-P7799	CD160	Mouse	Mammalian Cells	C-His
HY-P73499A	CD40	Human	Mammalian Cells	C-His
HY-P7819	CD207	Human	Mammalian Cells	N-His

Fc Receptor Proteins

Receptors of the fragment crystalizable (Fc) portion of immunoglobulin (Fc receptors, FcRs) are membrane molecules that are expressed on most innate and adaptive immune cells. FcRs belong to the immunoglobulin superfamily and interact with the Fc portion of antibodies to link humoral immune responses to cellular effector mechanisms. Receptors for all classes of immunoglobulins include IgG (FcγRI/CD64, FcγRII/CD32, FcγRIII/CD16, and FcRn), IgE (FCεRI, FcεRII), IgA (FcαRI/CD89), IgM (FcμR), IgD (FcδR) and IgA/IgM (Fcα/μR)[17][18].



Figure 4. Schematic illustration of Fc receptor interaction with an antibody-coated microbial pathogen[19]



Figure 5. Schematic figure of IgG[20]

Cat. No.	Product Name	Species	Source	Tag
HY-P70708	Fc gamma RIII/CD16	Mouse	Mammalian Cells	C-His
HY-P75209	Fc gamma RIIIA/CD16a	Human	Mammalian Cells	C-His
HY-P72657	FCAR/CD89	Human	Mammalian Cells	C-His
HY-P70711	Fc gamma RIIA/CD32a	Rat	Mammalian Cells	C-His
HY-P70669	CD64	Human	Mammalian Cells	C-His
HY-P70601	FCRN	Human	Mammalian Cells	C-His
HY-P72748	CD23/Fc epsilon RII	Human	Mammalian Cells	N-His
HY-P76800	Fc gamma RIIB/CD32b	Cynomolgus	Mammalian Cells	C-His
HY-P75204	Fc gamma RIIIB/CD16b	Human	*E.coli*	Biotinylated
HY-P72191	Fc epsilon RIA/FCER1A	Human	*E.coli*	His-SUMO
HY-P77363	FCAMR/CD351	Mouse	Mammalian Cells	C-His
HY-P70251	IgG3 Fc	Mouse	Mammalian Cells	Tag-free
HY-P72603	IgG2A Fc	Mouse	Mammalian Cells	Tag-free

Receptor Proteins

Receptors are protein molecules located on the cell surface or within the cytoplasm which are able to specifically recognize and bind to ligand molecules[21]. They are coupled to various signal transduction systems located both within the cell membrane and intracellularly, and can therefore regulate responses to the cellular/tissue microenvironment. Receptors are the molecular targets through which drugs produce their beneficial effects in various disease states[22][23].



Figure 6. Molecular signals underlying the anti-inflammatory effects of GLP1-GLP1R-based drugs[24]

Cat. No.	Product Name	Species	Source	Tag
HY-P70552	VEGFR-2	Human	Mammalian Cells	C-His
HY-P70714	HGFR	Human	Mammalian Cells	C-His
HY-P70352	GLP1R	Human	Mammalian Cells	C-Fc
HY-P70793	TrkB	Human	Mammalian Cells	C-His
HY-P7308	TrkA	Human	Mammalian Cells	Tag-free
HY-P70179	LIR-1/LILRB1	Human	Mammalian Cells	C-His
HY-P7485	Activin RIB/ALK-4	Human	Mammalian Cells	C-His
HY-P7467	AGER	Human	Mammalian Cells	C-His
HY-P71580	GFRAL	Human	*E.coli*	N-His-SUMO
HY-P72198	FSHR	Human	*E.coli*	N-His
HY-P70057	TYRO3/DTK	Mouse	Mammalian Cells	C-His

Enzymes & Regulators

Enzymes are biocatalysts found in biological systems that catalyze specific biochemical processes, and most enzymes are proteins. Many inherited human diseases, such as albinism and phenylketonuria, are caused by deficiencies in specific enzymes. The reactions catalyzed by enzymes can be inhibited by foreign or endogenous inhibitors. Many toxins and pharmacologically active substances act by inhibiting specific enzyme-catalyzed reactions[25][26][27].



Figure 7. Enzymatic catalysis[28]

Cat. No.	Product Name	Species	Source	Tag
HY-P7745	Cathepsin A	Human	Mammalian Cells	C-His
HY-P7442	ACE2	Human	Mammalian Cells	C-Fc
HY-P70351	MMP-9	Mouse	Mammalian Cells	C-His
HY-P70051	Aminopeptidase N/CD13	Mouse	Mammalian Cells	C-His
HY-P7474	ALDH1A1	Human	*E.coli*	N-His
HY-P7734	Carboxypeptidase B1/CPB1	Human	Mammalian Cells	C-His
HY-P70005	CTRB1	Human	Mammalian Cells	C-His
HY-P70010	CD73/5'-Nucleotidase	Human	Mammalian Cells	C-His
HY-P7452	ACOT13	Human	Mammalian Cells	C-His
HY-P70221	Acyl-protein thioesterase 2/LYPLA2	Human	*E.coli*	C-His
HY-P7479	Aldose 1-epimerase/GALM	Human	*E.coli*	C-His
HY-P72076	ALOX12	Human	*E.coli*	N-His
HY-P70260	Alpha-enolase/Enolase 1	Human	*E.coli*	C-His
HY-P73090	GSK-3 beta	Mouse	*E.coli*	N-His
HY-P7503	Angiogenin	Human	*E.coli*	Tag-free
HY-P73267	Kininogen-1	Mouse	Mammalian Cells	C-His
HY-P7602	Arginase-1	Human	Mammalian Cells	C-His

Complement System

Complement system, a complex network of plasma proteins that can be activated directly by invading pathogens or indirectly by pathogen-bound antibodies, plays a key role in host homeostasis, inflammation, and defense against pathogens. The complement system can be activated through three main pathways: the classical pathway (CP), the lectin pathway (LP), and the alternative pathway (AP). The initiation of these pathways depends on the final effector molecules: innocuous toxins (C4a/C3a/C5a), membrane attack complexes (MAC), and proteases (e.g. C3b)[29].



Figure 8. Complement System[30]

Cat. No.	Product Name	Species	Source	Tag
HY-P7863	Complement C3/C3a	Mouse	*E.coli*	Tag-free
HY-P7864	Complement C5/C5a	Human	*E.coli*	Tag-free
HY-P70031	CFHR2	Human	Mammalian Cells	C-His
HY-P70055	CFHR1	Human	Mammalian Cells	C-His
HY-P70102	CFHR5	Human	Mammalian Cells	C-His
HY-P7692	HABP1/C1QBP	Human	*E.coli*	C-His
HY-P71718	C1QA	Mouse	*P.pastoris*	N-His
HY-P71423	VSIG4	Human	Mammalian Cells	C-Fc
HY-P7890	Complement factor H/CFH	Human	Mammalian Cells	C-His
HY-P74371	C7/Complement component C7	Human	Mammalian Cells	C-His
HY-P74375	C2/Complement C2	Human	Mammalian Cells	C-His
HY-P74614	Protein S/PROS1	Human	Mammalian Cells	C-His
HY-P75464	C1QB	Human	Sf9 Insect Cells	C-His

Ubiquitin Related Proteins

Ubiquitin (Ub) is a small, highly conserved protein containing 76 amino acids that are ubiquitously expressed in eukaryotic cells. The covalent attachment of ubiquitin to target proteins is known as ubiquitination. Ubiquitination requires three distinct steps, which are ① activation of ubiquitin by Ub activase (E1), ② transfer of activated ubiquitin from E1 to the cysteine residues of Ub-binding enzyme (E2), and ③ attachment of ubiquitin to the lysine residues of the target protein by Ub ligase (E3)[31][32][33].



Figure 9. The ubiquitin (U)-protein conjugation cycle[34]

Cat. No.	Product Name	Species	Source	Tag
HY-P70843	NEDD8	Human	*E.coli*	Tag-free
HY-P701350	UBE2C	Human	Sf9 Insect Cells	N-His-Strep
HY-P71020	UBE2L6	Human	*E.coli*	C-His
HY-P71096	UCH-L1	Human	*E.coli*	C-His
HY-P71101	UBB	Human	*E.coli*	Tag-free
HY-P70974	SUMO2	Human	*E.coli*	N-His
HY-P7617	ATG3	Human	*E.coli*	Tag-free
HY-P71182	OTUB2	Human	*E.coli*	N-GST
HY-P71396	UBAP1	Human	*E.coli*	C-His
HY-P73537	XIAP	Human	*E.coli*	N-His
HY-P71643	SAE1	Human	*E.coli*	N-GST
HY-P70149	ISG15/UCRP	Human	*E.coli*	C-His
HY-P71395	UBA5	Human	*E.coli*	N-His
HY-P71402	UBE2H	Human	*E.coli*	N-GST
HY-P71407	UBE2R2	Human	*E.coli*	N-His

Viral Proteins

A complete infectious viral particle, called a virosome, consists of nucleic acid (DNA or RNA) and a capsid protein. The viral proteins of a mature assembled viral particle are known as viral structural proteins and may include core proteins of the core-shell (Gag proteins), enzymes packaged within the viral particle (Pol proteins), and membrane components (Env proteins). In addition, viral proteins include nonstructural proteins, regulatory proteins, and accessory proteins.



Figure 10. SARS-COV-2 structure[38]

Cat. No.	Product Name	Species	Source	Tag
HY-P7429	3C-like Proteinase	SARS-CoV-2	*E.coli*	N-His
HY-P7437	Nucleocapsid	SARS-CoV-2	*E.coli*	N-His
HY-P7436	S1 Protein	SARS-CoV-2	Mammalian Cells	Tag-free
HY-P70127	NSP1	SARS-CoV-2	*E.coli*	C-His
HY-P73290	Spike/S1	MERS-CoV	Sf9 insect Cells	C-His
HY-P70907	Envelope glycoprotein gp120	HIV-1	Mammalian Cells	C-His
HY-P74883	gp140	HIV-1	Mammalian Cells	C-Fc
HY-P70015	B18R	Vaccinia virus	Mammalian Cells	C-His
HY-P71478	Fusion glycoprotein F0/F	HRSVA	*E.coli*	N-His, B2M
HY-P73232	HA/Hemagglutinin	Influenza virus	Mammalian Cells	C-His
HY-P73239	NA/Neuraminidase	Influenza virus	Mammalian Cells	Tag-free
HY-P73533	Membrane protein	Zika virus	Mammalian Cells	C-Fc
HY-P73738	NS1 Protein	Dengue virus	Mammalian Cells	N-His
HY-P74188	E/Envelope Protein	West Nile Virus	*P.pastoris*	C-His
HY-P74354	Capsid protein	Hepatitis virus	*E.coli*	C-His

Biotinylated Proteins

Biotinylation is the process of covalently attaching biotin to a molecule, such as an amino acid. In general, biotinylation is rapid, specific, and unlikely to affect the natural function of the molecule due to the small size of biotin. Biotin is widely used in biomedical sciences due to its high affinity, rapid conductivity, and high specificity for binding to streptavidin/avidin. Importantly, biotinylated proteins are widely used as molecular tools in biotechnological applications[39].



Figure 11. Avi-tagged target protein biotinylated by BirA[40]

Cat. No.	Product Name	Species	Source	Tag
HY-P70546	PCSK9	Human	Mammalian Cells	C-His-HA-Avi
HY-P70721	PD-L1	Human	Mammalian Cells	C-Fc-Avi
HY-P70767	Siglec-15	Human	Mammalian Cells	C-Fc-Avi
HY-P70768	CD79B	Human	Mammalian Cells	C-His-Avi
HY-P70769	ACE2	Human	Mammalian Cells	C-His-Avi
HY-P70770	TROP-2	Human	Mammalian Cells	C-His-Avi
HY-P71175	NTNG1	Human	Mammalian Cells	C-Avi-His
HY-P71056	TGF beta 1	Human	Mammalian Cells	N-Avi
HY-P71421	VEGFR-2	Human	Mammalian Cells	C-His-Avi
HY-P73309	Neuropilin-1	Human	Mammalian Cells	C-His-Avi
HY-P73485	XIAP	Human	*E.coli*	N-Avi
HY-P78068	4-1BBR/TNFRSF9	Human	Mammalian Cells	C-hFc-Avi
HY-P72909	CD40	Human	Mammalian Cells	C-hFc-Avi
HY-P72885	Fc gamma RIIIA/CD16a	Human	Mammalian Cells	C-His-Avi
HY-P72343	BTN1A1	Human	Mammalian Cells	C-His-Avi
HY-P72389	LAG-3	Human	Mammalian Cells	C-His-Avi

GMP-grade Proteins

MedChemExpress GMP (Good Manufacturing Practice) recombinant proteins are manufactured under specific guidelines to ensure product quality and consistency. We offer GMP-grade recombinant proteins produced with quality documentation and full traceability, manufactured under independent QA oversight.

Cat. No.	Product Name	Species	Source	Tag
HY-P70637G	GMP TPO	Human	Mammalian cells	C-His
HY-P70757G	GMP SCF	Human	Mammalian cells	C-His
HY-P70454G	GMP IL-1 alpha	Human	*E.coli*	Tag-free
HY-P7044G	GMP IL-6	Human	*E.coli*	Tag-free
HY-P7032G	GMP IL-12	Human	Mammalian Cells	Tag-free
HY-P70760G	GMP IL-18	Human	Mammalian Cells	C-His
HY-P7038G	GMP IL-21	Human	*E.coli*	Tag-free
HY-P78551	GMP IL-2	Human	*E. coli*	Tag-free
HY-P7055G	GMP PDGF-BB	Human	*E.coli*	Tag-free

Animal-free Recombinant Proteins

Animal-free means that no animal-derived material is involved in the production process. Animal-free recombinant proteins have the same biological activity as proteins produced using standard laboratory techniques, while researchers do not have to worry at all about traces of animal components or mammalian pathogens affecting their experiments.

Cat. No.	Product Name	Species	Source	Tag
HY-P700017AF	Animal-Free BAFF	Human	*E.coli*	C-His
HY-P700018AF	Animal-Free beta-NGF	Human	*E.coli*	C-His
HY-P700020AF	Animal-Free BMP-11	Human	*E.coli*	C-His
HY-P700034AF	Animal-Free CCL2	Human	*E.coli*	N-His
HY-P700037AF	Animal-Free CD326	Human	*E.coli*	C-His
HY-P700042AF	Animal-Free CXCL11	Human	*E.coli*	N-His
HY-P700043AF	Animal-Free CXCL12	Human	*E.coli*	C-His
HY-P700045AF	Animal-Free CXCL2	Human	*E. coli*	N-His
HY-P700056AF	Animal-Free FGF-12	Human	*E.coli*	C-His
HY-P700066AF	Animal-Free FGF-4	Human	*E.coli*	C-His

Recombinant Protein Expression Service

- **One-stop service** from construct design to protein delivery
- Independently developed high expression vectors and strains
- Professional R & D team for recombinant protein
- Customized services of high-purity productions from milligram to gram scale
- Large-scale protein production platform



Service Types

MCE has five technical service platforms: E.coli expression system, yeast expression system, mammalian expression system, insect expression system and inclusion body protein renaturation.

Project	Content	Cycle	Deliverable
E.coli Expression System	Construction of Expression Vector	1-2 weeks	Expression Plasmid / Construction Report
	Protein Expression	1 week	Expression Report
	Flask Fermentation and Protein Purification	3 weeks	All Qualified Proteins / Purification Report
Inclusion Body Protein Renaturation	Construction of Expression Vector	1-2 weeks	Sequencing Report
	Protein Expression	1 week	Expression Report
	Small-scale Protein Renaturation	2-3 weeks	Refolding Protein
	Mass Protein Production	2-4 weeks	mg-g Level Refolding Protein
Yeast Expression System	Construction of Expression Vector	1-2 weeks	Construction Report
	Electrotransformation	2 weeks	5 Positive Strains
	Screening of High Copy Strains	1 week	1 High Copy Strain
	Protein Expression	3 weeks	Expression Report
	Protein Purification	3 weeks	Protein/Purification Report
Mammalian Expression System	Construction of Expression Vector	1-2 weeks	Construction Report
	HEK293/CHO Cell Transfection and Expression Detection	3 weeks	Expression Report
	1L HEK293/CHO Cell Transfection and Protein Purification	3 weeks	All Qualified Proteins / Purification Report
Insect Expression System	Construction of Expression Vector	2 weeks	Construction Report
	Preparation of Bacmid	1 week	Preparation Report
	Cell Transfection and Expression Detection	4-5 weeks	Expression Report
	Protein Purification	4 weeks	Protein / Purification Report

Customer Validation

Primary mouse pulmonary fibroblasts were treated with **TGF beta 1 Protein (HY-P7117)** to induce (myo) fibroblastic differentiation. The mRNA levels of PTPN13 and NOX4 in the (myo) fibroblasts were significantly increased after treatment, accompanied by increased expression of the (myo) fibroblast marker protein α-SMA[41].



Organoid beads were cultured with supplements of **Noggin (HY-P7086)**, **R-spondin 1 HY-P7114**, **FGF-4 (HY-P7014)**, **FGF-basic (HY-P7066)**, **SB-431542 (HY-10431)**, **Laduviglusib (HY-10182)**. The size, shape, and composition of the kidney organoids were highly reproducible[42].



IFN-γ induced the accumulation of IFI16. The expression of IFI16 in human lung adenocarcinoma epithelial cells (A549) treated with **IFN-γ (HY-P7025)** for 18 h was determined by western blot, and the intracellular localization of IFI16 in A549 cells treated with IFN-γ for 12 h was determined[43].



Selected Publications Citing Use of MCE Recombinant Proteins

Publications	Product Name	Cat. No.
Cell. 2024 Sep 18:S0092-8674(24)00977-2.	EGF	HY-P7109
Cell. 2024 Sep 4:S0092-8674(24)00904-8.	BMP-2	HY-P7006
Cell. 2024 Apr 25;187(9):2288-2304.e27.	IL-10; IL-2; IL-6	HY-P7030; HY-P7037; HY-P7044
Cell Host Microbe. 2023 Nov 8;31(11):1820-1836.e10.	IFN-beta	HY-P73130
Nat Chem. 2024 Nov;16(11):1811-1821.	PCBP1	HY-P74649
ACS Nano. 2024 Oct 1;18(39):26784-26798.	MMP-2	HY-P73810
Nat Commun. 2024 Sep 12;15(1):7898.	Myeloperoxidase	HY-P70282
Nat Commun. 2024 Sep 12;15(1):7996.	FGF basic	HY-P7004
Ann Rheum Dis. 2024 Sep 4:ard-2024-225565.	NCL	HY-P71760
Nat Cell Biol. 2024 Sep;26(9):1597-1612.	IgG1	HY-P73143
Adv Sci (Weinh). 2024 Aug 5:e2404408.	HMGB1	HY-P70570
Adv Sci (Weinh). 2024 Aug 8:e2406633.	IL-18	HY-P70591
Nat Commun. 2024 Aug 5;15(1):6626.	UBE2D3; STUB1	HY-P70998; HY-P71340
Nat Commun. 2024 Aug 16;15(1):7077.	GMP IL-4; M-CSF	HY-P70750G; HY-P7085A
ACS Nano. 2024 Aug 20;18(33):22194-22207.	MMP-9	HY-P73807
Adv Sci (Weinh). 2024 Jul 1:e2403640.	Prolactin	HY-P700261
Nat Commun. 2024 Jul 27;15(1):6344.	FGF basic; Betacellulin	HY-P7004; HY-P7005
Mol Cancer. 2024 Jul 9;23(1):141.	EGF	HY-P7109
Adv Sci (Weinh). 2024 Jul 26:e2404069.	IL-4	HY-P70653
ACS Nano. 2024 Jul 2;18(26):17119-17134.	GM-CSF	HY-P7361
Adv Sci (Weinh). 2024 Jun 18:e2400115.	YES1	HY-P73486
Adv Mater. 2024 Jun 17:e2402738.	Osteopontin	HY-P71786
J Extracell Vesicles. 2024 Jun;13(6):e12467.	CD81	HY-P74264
Acta Pharm Sin B. 2024;14(10):4360e4377.	IL-12	HY-P70666
Adv Sci (Weinh). 2024 Jun 20:e2404171.	TNF-alpha	HY-P7058
Adv Funct Mater. 2024:2404549.	M-CSF; RANK L	HY-P7085; HY-P7425
Adv Mater. 2024 May;36(19):e2311964.	Osteopontin	HY-P78358
Cancer Cell. 2024 Apr 8;42(4):682-700.e12.	CHI3L1; MMP-2	HY-P70030; HY-P70268
Cancer Commun (Lond). 2024 Apr;44(4):469-490.	FGF basic; EGF	HY-P7004; HY-P7109
Adv Funct Mater. 2024 Apr;34(14):2310416.	GDF-5	HY-P72633
Adv Mater. 2024 Mar 24:e2401384.	IL-4	HY-P7080
Immunity. 2024 Mar 12;57(3):495-512.e11.	IL-1 beta; IL-1RA	HY-P7073; HY-P72566
Nat Genet. 2024 Feb;56(2):294-305.	EGF	HY-P7109

References:

[1] Jun-Ming Zhang. Cytokines, inflammation, and pain. Int Anesthesiol Clin. 2007 Spring;45(2):27-37.

[2] Pedro Berraondo, et al. Cytokines in clinical cancer immunotherapy. Br J Cancer. 2019 Jan;120(1):6-15.

[3] Ferreira, Vinicius, et al. Cytokines and Interferons: Types and Functions. In Autoantibodies and Cytokines, edited by Wahid Khan. London: IntechOpen, 2018.

[4] Xiaochen Ren, et al. Growth Factor Engineering Strategies for Regenerative Medicine Applications. Front Bioeng Biotechnol. 2019; 7: 469.

[5] Stone WL, Leavitt L, Varacallo M. Physiology, Growth Factor. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022 Jan.

[6] Ozlem Guzeloglu-Kayisli, Hugh S Taylor, et al. The role of growth factors and cytokines during implantation: endocrine and paracrine interactions. Semin Reprod Med. 2009 Jan;27(1):62-79.

[7] Yaping Zhang, et al. Functions of Immune Checkpoint Molecules Beyond Immune Evasion. Adv Exp Med Biol. 2020;1248:201-226.

[8] Xing He, et al. Immune checkpoint signaling and cancer immunotherapy. Cell Research. 2020;30:660-669.

[9] Marin-Acevedo JA, et al. Next generation of immune checkpoint therapy in cancer: new developmentsand challenges. J Hematol Oncol. 2018;11(1):39.

[10] Christopher DeRenzo, et al. Genetic Modification Strategies to Enhance CAR T Cell Persistence for Patients With Solid Tumors. Front Immunol. 2019 Feb 15;10:218.

[11] M Essand, et al.Genetically engineered T cells for the treatment of cancer. J Intern Med. 2013 Feb;273(2):166-81.

[12] Zhenguang Wang, et al. New development in CAR-T cell therapy. J Hematol Oncol. 2017; 10: 53.

[13] Lekha Mikkilineni, et al.Chimeric antigen receptor T-cell therapies for multiple myeloma. Blood. 2017 Dec 14;130(24):2594-2602.

[14] Julius M. Cruse, Robert E. Lewis, and Huan Wang. CLUSTER OF DIFFERENTIATION (CD) ANTIGENS. Immunology Guidebook. 2004 : 47-124.

[15] Tomas Kalina, et al. CD Maps-Dynamic Profiling of CD1-CD100 Surface Expression on Human Leukocyte and [16] Lymphocyte Subsets. Front Immunol. 2019 Oct 23;10:2434.

[16] Dario Gosmann, Angela M. Krackhardt, Calogero D'Alessandria, et al. P. romise and challenges of clinical non-invasive T-cell tracking in the era of cancer immunotherapy. EJNMMI Res. 2022 Jan 31;12 (1):5.

[17] Maree S. et al. Fc Receptors. Adv Exp Med Biol. 2008;640:22-34.

[18] Sanae Ben Mkaddem, et al. Understanding Fc Receptor Involvement in Inflammatory Diseases: From Mechanisms to New Therapeutic Tools. Front Immunol. 2019 Apr 12;10:811.

[19] D. Mancardi. Fc Receptor-Dependent Immunity. Reference Module in Biomedical Sciences. 2014 : B978-0-12-801238-3.00119-7.

[20] Mårtensson, A. Development of an Antigen-independent Affinity Assay to Study the Binding of IgG to Fc Gamma Receptors. 2012.

[21] S. Xiong, et al. Human Immune System,Editor(s): Anders rahme,Comprehensive Biomedical Physics,Elsevier,2014,Pages 91-114.

[22] Carl-Henrik Heldin, et al. Signals and Receptors. Cold Spring Harb Perspect Biol. 2016 Apr; 8(4): a005900.

[23] Rita Santos, et al. A comprehensive map of molecular drug targets. Nat Rev Drug Discov. 2017 Jan; 16(1): 19-34.

[24] Young-Sun Lee, et al. Anti-Inflammatory Effects of GLP-1-Based Therapies beyond Glucose Control. Mediators Inflamm. 2016;2016:3094642.

[25] Peter K. et al. Enzymes: principles and biotechnological applications. Essays Biochem. 2015 Nov 15; 59: 1-41.

[26] Antonio Blanco, et al. Enzymes, Editor(s): Antonio Blanco, Gustavo lanco,Medical Biochemistry,Academic Press,2017,Pages 153-175.

[27] Cooper GM. The Cell: A Molecular Approach. 2nd edition. Sunderland (MA): Sinauer Associates; 2000.

[28] Liubov Poshyvailo. Modelling and simulations of enzyme-catalyzed reactions. National University of "Kyiv-Mohyla Academy" Faculty of Natural Sciences, Department of Physics and Mathematics.2015.

[29] Janeway CA Jr, et al. The complement system and innate immunity. Immunobiology: The Immune System in Health and Disease. 5th edition. New York: Garland Science; 2001.

[30] Guillermina Girardi, et al. Essential Role of Complement in Pregnancy: From Implantation to Parturition and Beyond. Front Immunol. 2020; 11: 1681.

[31] Lan K Nguyen, et al. When ubiquitination meets phosphorylation: a systems biology perspective of EGFR/MAPK signaling. Cell Commun Signal. 2013 Jul 31;11:52.

[32] Guo HJ, et al. Biochemistry, Ubiquitination. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022 Jan.

[33] Fu Shang, et al. Ubiquitin-proteasome pathway and cellular responses to oxidative stress. Free Radic Biol Med. 2011 Jul 1; 51(1): 5-16.

[34] Hochstrasser, M. Origin and function of ubiquitin-like proteins. Nature. 2009 Mar 26;458(7237):422-9.

[35] Viral Structural Proteins. NCBI. Literature. MeSH Database. Year introduced: 1990.

[36] Jennifer Louten. Virus Structure and Classification. Essential Human Virology. 2016 : 19-29.

[37] Gelderblom HR. Structure and Classification of Viruses. In: Baron S, editor. Medical Microbiology. 4th edition. Galveston (TX): University of Texas Medical Branch at Galveston; 1996. Chapter 41.

[38] Chandra Mohan, et al. A Brief Study on Rate of Morphological and Genomic Similarities between SARS and COVID-19. Int J Pharma Res Health Sci. 2020; 8 (3): 3167-71.

[39] Granhøj, J., et al. A bacterial display system for effective selection of protein-biotin ligase BirA variants with novel peptide specificity. Sci Rep. 2019; 9, 4118.

[40] Huijun Xue, et al. Utilizing Biotinylated Proteins Expressed in Yeast to Visualize DNA–Protein Interactions at the Single-Molecule Level. Front Microbiol. 2017; 8: 2062.

[41] Jiwei Hou, et al. Co-delivery of siPTPN13 and siNOX4 via (myo)fibroblast-targeting polymeric micelles for idiopathic pulmonary fibrosis therapy. Theranostics. 2021 Jan 9;11(7):3244-3261.

[42] Chengyong He, et al. Black Phosphorus Quantum Dots Cause Nephrotoxicity in Organoids, Mice, and Human Cells. Small. 2020 Jun;16(22):e2001371.

[43] Zhimin Jiang, et al. IFI16 directly senses viral RNA and enhances RIG-I transcription and activation to restrict influenza virus infection. Nat Microbiol. 2021 Jul;6(7):932-945.

Inhibitors • Screening Libraries • Proteins

www.MedChemExpress.com

MedChemExpress USA

Tel: 609-228-6898 **E-mail:** sales@MedChemExpress.com

Fax: 609-228-5909 **Tech Support:** tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite F, Monmouth Junction, NJ 08852, USA

MedChemExpress Europe

Tel: +4686500910 **E-mail:** eu.sales@MedChemExpress.com

Address: Bergkällavägen 37C 192 79 Sollentuna SWEDEN

For research use only.

We do not sell to patients.

Master of

Bioactive Molecules

www.MedChemExpress.com